

Review article

A decade of plant proteomics in South Korea: the International plant proteomics organization (INPPO) perspective and involvement

Su-ji Lee¹, Kyu Young Kang^{2,3}, Nam-Soo Jwa⁴, Dea-Wook Kim⁵, Ganesh Kumar Agrawal⁶, Abhijit Sarkar^{6,7}, Renu Deswal⁸, Jenny Renaut⁹, Dominique Job¹⁰, Randeep Rakwal^{6,11,12*}, and Sun Tae Kim^{1*}

¹Department of Plant Bioscience, Pusan National University, Miryang, 627-706, South Korea

²Division of Applied Life Science (BK21 program), Gyeongsang National University, Jinju 660-701, South Korea

³Plant Molecular Biology and Biotechnology Research Center, Gyeongsang National University, Jinju 660-701, South Korea

⁴Department of Molecular Biology, Sejong University, Gunja-dong, Seoul 143-747, South Korea

⁵National Institute of Crop Science, Rural Developmental Administration, Suwon 441-857, South Korea

⁶Research Laboratory for Biotechnology and Biochemistry (RLABB), GPO Box 13265, Kathmandu, Nepal

⁷Department of Botany, Banaras Hindu University, Varanasi 221005, India

⁸Department of Botany, University of Delhi, Delhi-7, India

⁹Centre de Recherche Public-Gabriel Lippmann, Department of Environment and Agrobiotechnologies, Belvaux, GD, Luxembourg

¹⁰CNRS/UCBL/INSA/Bayer CropScience Joint Laboratory, UMR 5240, Bayer CropScience, 14-20 rue Pierre BAIZET, F-69263, Lyon cedex, France

¹¹Graduate School of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba Ibaraki 305-8572, Japan

¹²Department of Anatomy I, Showa University School of Medicine, 1-5-8 Hatanodai, Shinagawa, Tokyo 142-8555, Japan

*Corresponding author: Sun Tae Kim, stkim71@pusan.ac.kr; Randeep Rakwal, plantproteomics@gmail.com

Abstract

In this review, we discuss the contribution of proteomics focusing on plant responses to various environmental stimuli in South Korea. Due to improvements in proteomics methods and applications in various research fields in South Korea, plant proteomics have and will continue to provide systematic approaches to address biological questions and understanding of the network between plants and environmental stimuli. We also introduce the International Plant Proteomics Organization (INPPO; www.inppo.com), a global initiative to advance plant proteomics research worldwide that is also designed to highlight national plant proteomics topics via a common global platform. Thus, we expect INPPO to generate further interest and activity among South Korean plant proteomics researchers and within the plant biology scientific community. Finally, to help network the plant proteomics community in South Korea, the creation of an INPPO-South Korea chapter is proposed.

Keywords: Abiotic stress, Biotic stress, Environment, INPPO, Plant Proteomics.

Abbreviations: HSPs: heat shock proteins; MS: mass spectrometry; LCM: liquid culture medium; PEG: polyethylene glycol; PCD: programmed cell death; PVDF: polyvinylidene difluoride; RuBisCO: ribulose-1,5-bisphosphate carboxylase/oxygenase; 2-DGE: two-dimensional gel electrophoresis; SCCs: suspension cultured cells.

Introduction***Crop plants and the start of proteomics and technological advances***

Crops and plants of economical importance are the primary targets of proteomics research in South Korea, which is generally conducted to improve seed quality and yield. Rice, being a major crop, has always been a major focus of research. South Korea has large quantities of natural and cultivated rice germplasm and researchers that have been thoroughly trained in investigation of these resources. In the past decade, proteomics has increasingly been applied as a high-throughput technology to complete our understanding of plant systems under diverse environments (Thiellement et al., 2007, Agrawal and Rakwal 2008a). Plant proteomics in South Korea is believed to have started in the late 1990s, when a research

group led by Professor Kyu Young Kang initiated proteomics investigations of rice plants at Gyeongsang National University. In their first published study, they described a novel method for depletion of the most abundant leaf protein in rice, ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO), via a simple polyethylene glycol (PEG) fractionation technique (Kim et al., 2001). This pioneering method demonstrated how a simple extraction method combined with two-dimensional gel electrophoresis (2-DGE) could increase the detection of rare or low-abundant proteins, and hence tremendously increase coverage of the leaf proteome. The same group also developed a 1-DGE (SDS-PAGE; sodium dodecyl sulfate polyacrylamide gel electrophoresis)-based methodology to fractionate and enrich high-/low-molecular mass proteins of the rice leaf (Kim et al., 2003a). In that study, the ability to combine 1-DGE with

2-DGE was also demonstrated. This method revealed the usefulness of a classical SDS-PAGE in the proteomics field for in-depth investigation of a proteome. In later years, successive development of several methodologies in rice proteomics allowed Kyu Young Kang's group to conduct thorough proteomics investigations of rice and develop comprehensive maps of the proteome responses of rice to multiple biotic and abiotic stresses (reviewed in Agrawal and Rakwal 2006, 2008b, 2011, Agrawal et al., 2009, 2010). These and other technical advancements, along with the confidence in proteomics technology for addressing biological questions (Bradshaw and Burlingame 2005) have led to its widespread utilization by a number of research groups at Universities and Institutions. Most, if not all of these groups are highlighted onto the map of South Korea in Fig. 1.

Plant proteomics database in South Korea

Increases in the number of research groups and publications reflect the rapid progress in plant proteomics that has occurred in South Korea in the past decade. Many scientists are now working intensively on numerous biological questions pertaining to agriculture crops and plants of their interest (Fig. 2). The plant systems that are being used for such investigations range from the dicot model *Arabidopsis thaliana* to the monocot cereal model crop rice (*Oryza sativa* L.), and include numerous other important crop plants, such as wheat, maize, soybean, rapeseed (canola), buckwheat, and banana. Investigations of plant responses to biotic (i.e., bacterial, fungus, and viral pathogens) and abiotic (i.e., agrochemicals, cold, drought, heat, heavy metals, light, ozone, salt, UV-B, volatile organic compounds, and water logging) stresses through proteomics techniques have been the major focus of South Korean plant proteomers since the technique was first introduced. In fact, this tremendous progress in plant proteomics was highlighted in a special issue of the Journal of Plant Biotechnology (a peer reviewed South Korean Scientific Journal published by the Korean Society of Plant Biotechnology) (Lee et al., 2011). This special issue in a national journal, and increased numbers of research publications to keep up with leading international journals, advocated the achievements made in South Korean plant proteomics and increased the awareness of the potential for the use of proteomics to study plants among plant biologists. Such attention and increased funding are required to induce the next green revolution to feed and meet the demands of the ever-growing human population under global climate changes (Swaminathan 2000). In the following sections, we look at two critical aspects of plant proteomics research in South Korea, abiotic and biotic stress proteomics in plants, particularly crops such as rice.

Abiotic stress proteomics

Rice proteomics has achieved remarkable progress in the use of high-throughput techniques, as well as generation of meaningful biological data pertaining to developmental stages, tissues and organs, and against biotic and abiotic stresses. Agrawal and Rakwal have systematically and comprehensively reviewed the progress in rice proteomics during 2000 to 2010 (Agrawal and Rakwal 2006, 2008b, 2011, Agrawal et al., 2009, 2010, Rakwal and Agrawal 2003). In Korea, a total of 29 papers on plant proteome responses to abiotic stresses including agrochemicals, heavy metals, cold, heat, drought, water logging, and salt stresses were published from 2005 to 2010. These papers are summarized in Fig. 3, along with an illustration of the proteomics approaches. Most experiments to




Fig 1. Plant proteomics research in South Korea and a potential INPPO-South Korea platform.

investigate abiotic stresses responses were carried out using leaf and root tissues of rice and *Arabidopsis*. Below, we briefly summarize proteomics studies of abiotic stresses in rice as a model crop system. These studies have increased our knowledge of abiotic stress-responsive proteins and regulatory pathways and mechanisms in rice.

Agrochemical and heavy metal stress

A total of 11 papers published between 2005 and 2010 have analyzed changes in the proteome to investigate the effects of two agrochemicals, glyphosate and glufosinate, and of heavy metals such as arsenic (As), cadmium (Cd), and copper (Cu). Proteome-wide changes in response to heavy metals comprise the largest set of papers (five papers), with As in leaf (Lee et al., 2008, Ahsan et al., 2010) and root (Ahsan et al., 2008a), Cd in leaf/root (Lee et al., 2010) and seedling (Ahsan et al., 2007a), and Cu in seedling (Ahsan et al., 2007b). Proteomics analysis revealed that most of the proteins related to energy production and metabolism, such as the RuBisCO large subunit (LSU) and detoxification, were increased. Conversely, proteomics analysis of rice leaves treated with glyphosate (an herbicide used to inhibit aromatic amino acid synthesis) and glufosinate (an herbicide inhibiting glutamine synthetase to catalyze ammonia assimilation) revealed that antioxidant enzymes and photosynthetic/metabolic-related proteins such as RuBisCO LSU were decreased (Ahsan et al., 2008b, Lee et al., 2008), suggesting that plant cells maintain a homeostasis in photosynthetic and antioxidative machineries corresponding to the environmental stresses imposed on plants.

Temperature stress

In addition, Lee et al. (2007a, 2009) investigated cold-responsive proteins in leaves and roots of rice exposed to chilling stress conditions (5 or 10°C) in a time-course experiment. Proteins such as cysteine proteinase, thioredoxin peroxidase, a RING zinc finger protein-like, drought-inducible late embryogenesis abundant protein, and a fibrillin-like protein were identified. Another study of the leaf proteome of rice exposed to 42°C for 12 and 24 h was also conducted (Lee et al., 2007b). Based on increased biochemical changes such as ion leakage and lipid peroxidation, it was suggested that high temperature treatment induced oxidative stress in rice leaves.




Fig 2. Studies, objectives, and contributions of the plant proteomics in Korea (adopted from Lee et al., 2011).




Fig 3. A summary of published papers on rice abiotic stress proteomics (adopted from Lee et al., 2011).

Seventy-three out of 1000 proteins separated by 2-DGE were identified by mass spectrometry (MS), 48 of which were identified as heat shock proteins (HSPs), energy and metabolism-, redox homeostasis-, and regulation-related proteins after heat treatment. One important finding was that many small HSPs (sHSPs) are newly up-regulated. The sHSPs play a central role in the complex cellular network (Wehmeyer et al., 1996), and it has been suggested that plants protect themselves from heat stress using a complex mechanism, of which sHSPs are a part (Lee et al., 2007b).

Biotic stress proteomics

Biotic stressors such as bacteria, viruses, fungi, insects, and weeds not only have the potential to cause damage, but also are the direct cause of vast economic losses of crops. Proteomics studies against biotic stress factors have been conducted in *Arabidopsis*, pepper, rice, and wheat (Fig. 4). Many of these studies were conducted using the blast lesion mimic (*blm*) mutant. This mutant is associated with biotic stress response proteins coupled to pathogens or defense-related signals. As a major fungal pathogen, an interaction between *Magnaporthe*

oryzae (*M. oryzae*) causing rice blast disease and rice plants was reported by Jung et al., (2005, 2006), Kim et al., (2009), and Ryu et al., (2009). We discuss these investigations in the following sections while focusing on studies dealing with the interaction of *M. oryzae* and rice.

Rice plant-pathogen interactions and programmed cell death

M. oryzae and rice

The rice blast fungus *M. oryzae* is a causal agent of rice blast disease, one of the most serious pathogens of the rice plant in most rice-growing regions of the world. Kyu Young Kang and colleagues have meticulously carried out comparative proteomics analyses of the interactions between *M. oryzae* and rice (Fig. 5). The first proteomics analysis was conducted in 2003 using a rice suspension cultured cells (SCCs) system (Kim et al., 2003b) inoculated with rice blast fungus or treated with elicitors to identify pathogen-responsive proteins. In 2004, these authors extended the study to characterization of *M. oryzae*-responsive proteins from rice leaves infected with




Fig 4. A summary of published papers on rice biotic stress proteomics (adopted from Lee et al., 2011).




Fig 5. History of rice proteomics during rice-blast pathogen interaction.

compatible (KJ301) and incompatible (KJ401) fungus (Kim et al., 2004a). The use of the PEG (15% w/v) fractionation technique previously developed to remove RuBisCO into the pellet fraction (Kim et al., 2001) enabled a comprehensive analysis of the *M. oryzae*-responsive proteins from rice (cv. Jinheung) leaves. Among proteins identified by N-terminal sequencing and MALDI-TOF-MS were the probenazole-induced protein (PBZ1), OsRLK, and OsPR10, which were highly upregulated in both SCCs and leaves in response to *M. oryzae* attack. These results also revealed that the expression of PBZ1, OsRLK, and OsPR10 was faster and higher in the incompatible interactions than compatible interactions. These results imply that rapid and high-level expression of these genes may provide host plants with a self-defense mechanism against pathogen attack. Accordingly, these proteins might also serve as potential defense-related biomarkers in rice.

PBZ1 and programmed cell death

PBZ1, which is a PR 10 family protein, was found to be induced in both rice SCCs and leaves inoculated with blast fungus (Kim et al., 2003b, 2004a). Despite the above research linking PBZ1 with defense/stress response, its biological function remained unknown. Interestingly, Kim and coworkers (Kim et al., 2008) explored the role of PBZ1 as a putative cell death marker in rice. Based on these studies, PBZ1 was found to be highly accumulated in tissues undergoing programmed cell death (PCD). The correlation between the accumulation of PBZ1 and PCD was demonstrated during leaf senescence, root aerenchyma formation, coleoptiles senescence, and root cap and seed aleurone layer formation (Kim et al., 2008) by monitoring PBZ1 expression level using Western blotting, immunohistochemical analysis, and promoter analysis. The results showed that the expression levels and localizations of

PBZ1 were tightly correlated with plant cell death. These results strongly suggest that PBZ1 is either a molecular marker in rice defense responses or can serve as a novel potential biomarker for cell death/PCD in rice.

M. oryzae and rice, and their secretomes

Secretomes are an emerging area for understanding interactomes between hosts and pathogens in the extracellular space within the rice leaf, which serves as a front line of defense against invading blast fungus (see review, Agrawal et al., 2010). To the best of our knowledge, there have been no reports of the *in vitro* and *in vivo* secretomes of rice leaves infected with compatible and incompatible races of *M. oryzae*. Previously, secretome analysis of rice SCCs treated with *M. oryzae* or its elicitors resulted in the identification of 21 differential protein spots (Kim et al., 2009). Mass spectrometry analysis resulted in the identification of nine chitinases, expansin, five 33 kDa secretory proteins (DUF 26), and two germin/oxalate oxidases protein. One of the identified chitinase proteins revealed strong enzymatic activity in an in-gel assay. Corresponding gene expression patterns were also validated by RT-PCR, which revealed a much earlier induction of chitinase, DUF26, and expansin in incompatible rather than compatible interactions. This secretome analysis by Kim et al. was the first to provide information about the mechanism of rice defense response against *M. oryzae* and its elicitor. However, the *in vivo* secretome of any plant-pathogen interaction has yet to be systematically investigated. In the near future, the established *in vivo* rice apoplastic proteome will provide important information enabling better understanding of the rice-*M. oryzae* interaction. While studying the blast-fungus interaction proteome, it is also desirable to examine the pathogen portion of the secretome to obtain better insight into the rice self-defense mechanism. During the host infection process, *M. oryzae* differentiates into a special structure, known as an appressorium, to penetrate the leaf surface, followed by invasive hyphae in an early stage of the biotrophic infection phase. Kim et al., (2004b) conducted a study to identify the proteins induced in the rice blast fungus *M. oryzae* during appressorium formation using 2-DGE. Protein spots differentially expressed during appressorium formation were confirmed from gels after 2-DE analysis of proteins labeled with ³⁵S methionine and stained with silver nitrate. This allowed isolation and identification of three proteins (20S proteasome a subunits, scytalone dehydratase, and serine carboxypeptidase Y) that were differentially accumulated during appressorium formation. In particular, the expression levels of the 20S proteasome proteins identified by Western blot analysis during appressorium formation and nutrient starvation increased greatly. Thus, the function of the 20S proteasome proteins might provide important information pertaining to the cellular mechanism involved in appressorium formation. Wang et al. (2011) recently presented a comparative 2-DGE-based proteomics approach that revealed the secretome of *M. oryzae* responsive to N starvation using cultured liquid complete medium, minimal medium, and N starvation minimal medium. A total of 85 protein spots differentially responsive to N starvation were identified by MALDI-TOF-MS and μ LC-ESI-MS/MS. The identified proteins were mainly cell wall hydrolase enzymes (22.4%), protein and lipid hydrolases (24.7%), ROS detoxifying proteins (22.4%), and proteins with unknown function (14.1%). The results showed that several of the identified secreted proteins had previously been found to be associated with fungal growth and pathogenicity.

Furthermore, semi-quantitative RT-PCR analysis of N-responsive secreted proteins revealed a good correlation between RNA and protein levels, suggesting that they might also provide insight into *M. oryzae* pathogenicity and development of methods to control infection during the early stages. Most recently, Jwa et al. conducted secretome analysis of *M. oryzae* by mimicking the early stages of infection *in vitro* using a glass plate (GP), polyvinylidene difluoride (PVDF) membrane, and liquid culture medium (LCM) (Jung et al., 2012). Specifically, they used 2-DGE and tandem mass spectrometry analyses to identify 53 non-redundant proteins including previously known and novel secreted proteins. Most of the proteins were found to be involved in cell wall modification, ROS detoxification, lipid modification, metabolism, and protein modification categories. This unique study employing both hydrophobic (GP and PVDF) and hydrophilic (LCM) conditions provided a survey of the *M. oryzae* secretome that can be used to further our understanding of the early interactions between *M. oryzae* and rice leaves. In conclusion, many proteins have been found to be differentially expressed in either the plant or pathogens during their interactions. However, it is likely that information providing a clear understanding of biological processes occurring during rice and *M. oryzae* interaction will require further investigations. More in-depth analyses of the special and temporal distribution of responding proteins (e.g., *in-planta* apoplastic secretome changes) will help elucidate details pertaining to pathogen invasion strategies. In the technical aspect for analysis of proteomics, the 2-DGE approach has its own limitations in identifying extreme proteins such as highly basic, acidic, or membrane proteins as well as proteins with low abundance. Use of complementary proteomics approaches such as shotgun coupled with 1-DGE using MudPIT and iTRAQ (Weckwerth 2011) will be helpful in expanding and saturating the defense responsive proteins during plant and microbe interactions.

Future challenges for plant proteomics research in South Korea

Despite considerable progress in plant proteomics research in South Korea, there are numerous challenges ahead that must be met before the potential of proteomics technology in plant science is fully realized. We would like to bring out several points for discussion among various research groups. These are to: i) have an open sharing of information, such as proteomics techniques (i.e., extraction protocols, 1-DGE, 2-DGE, peptide identification, and protein assignment), without overlapping the objectives of each working group; ii) address major ground level problems in the country; and iii) help translate the outcome of the research to the field to produce next-generation crops with actively involving agronomists and breeders. With regard to the production of stress-resistant crops (point iii) it is without doubt that proteomics data pertaining to the complexity of plant responses to biotic and abiotic environmental factors can facilitate selection of potential biomarkers of plant tolerance to these deleterious stresses. Such biomarkers could be useful for plant biologists and breeders in marker-assisted crop and plant breeding. However, it is still necessary to determine a method of translating this information for the breeders. The National Institute of Crop Science (NICS) under the Rural Development Administration (RDA) in South Korea faces a similar problem. Breeders at NICS ask how the information accumulated from proteomics studies will facilitate sustainable crop production under unfavourable environmental

conditions or provide tolerance or resistance to crops under biotic and abiotic stresses. Accordingly, it is important to convey that, although plant genomes contain about 25-50,000 genes that are estimated to encode more than 1,000,000 protein isoforms because of the existence of extensive post-translational modifications of proteins, it is anticipated that such modifications (e.g., phosphorylations, methylations, acetylations, prenylations...) would constitute biomarkers of plant stress response, which cannot be identified by classical genetic approaches. Therefore these protein markers will comprise the next generation of genetic markers traditionally used in breeding, allowing further progress in plant improvement. With regards to the concerns of breeders, it is essential to tap the resources pertaining to crop varieties that are being generated in institutes such as the NICS to help bridge the gap and expectations of plant breeders by making direct links with the proteomics specialists in universities in South Korea.

The INPPO initiative and its association with plant proteomics

The International Plant Proteomics Organization (INPPO; www.inppo.com) has been founded with a major goal of the global sharing of knowledge databases gathered by plant proteomics studies worldwide. One of the major objectives of INPPO is to bring national plant proteomics issues out in the open via a common global platform. A global platform has always helped promote science and technology, and their application in human health and food security. INPPO has been established as a global platform for plant proteomics that includes ten initiatives (for details, see INPPO viewpoint paper, Agrawal et al., 2011; and INPPO Highlights, Agrawal et al., 2012b). The objectives of INPPO are to intensify cooperation in the field of plant proteomics, establish complete plant proteomes, carry out comparative and translational proteomics (Agrawal et al., 2012a), establish centralized databases, organize conferences and workshops, integrate proteomics-related activities and disseminate them through the INPPO website, provide education and training, develop interactions with other proteomic initiatives such as HUPO (the Human Proteome Organization) and MASCP (the Multinational *Arabidopsis* Steering Committee for Proteomics), and help address biological and societal questions.

To continue in line with these INPPO initiatives, we envision a national Plant Proteomics Chapter in South Korea under the INPPO umbrella, which we propose as the 'INPPO – South Korean Plant Proteomics Chapter (INPPO – South Korea)'. We sincerely hope that such a forum at the national level will bring together all plant proteomers, biologists, and breeders at a common platform to discuss the pressing problems faced, in particular by the plant and agricultural scientific communities. We look forward to the active participation of the plant proteomics community in this endeavor. Moreover, to help the younger plant proteomers progress to the next level in proteomics research, high-level training under the guidance of experts in national and international laboratories with specific expertise will be necessary. Initially, available educational infrastructure in the Universities will have to include basic proteomics facilities to generate interest in and train undergraduates and graduates in the latest proteomics skills. In the long-term, creation of more state-of-the-art high-throughput proteomics facilities dedicated to plant proteomics will be required. One good example is the facility for proteomics at Yonsei University in Seoul, which is a hub for Korea HUPO

that primarily focuses on mammalian systems, but also includes plant proteomers. However, to accomplish such a large jump in plant proteomics research, one or two more similar centers will have to be created, depending on the collective will and use of the plant sciences community. Only then can the potential of the younger generation of Korean plant proteomers be fully realized. Finally, in keeping with the INPPO spirit of cohesion, coordination, and synergy, we look forward to support from our colleagues in South Korea in taking plant proteomics research to the next level.

Conclusion

In this review, we highlight proteomics research cases and trends focusing on plant responses to various stimuli including abiotic and biotic stresses in Korea. Furthermore, this review provides advanced researchers with information about the “who and who” of Korean plant proteomics dealing with plant response and networking to its environment. With the opening of the INPPO-South Korea plant proteomics chapter, we not only hope to see plant proteomics gain prominence nationally but also set an example for other countries to network plant proteomers.

Acknowledgements

We thank the plant scientists and the proteomics community in South Korea for their research and cooperation. The authors also acknowledge the INPPO platform for this initiative, which led to collaboration by scientists of different disciplines for this review. This work was supported by a grant from the Next-Generation BioGreen 21 Program (SSAC, grant#: PJ008107), Rural Development Administration (RDA), Republic of Korea.

References

- Agrawal GK, Job D, Zivy M, Agrawal VP, Bradshaw RA, Dunn MJ, Haynes PA, van Wijk KJ, Kikuchi S, Renaut J, Weckwerth W, and Rakwal R (2011) Time to articulate a vision for the future of plant proteomics – a global perspective. An initiative for establishing the International Plant Proteomics Organization (INPPO). *Proteomics* 11:1559-1568.
- Agrawal GK, Jwa NS, Lebrun M-H, Job D, and Rakwal R (2010) Plant secretome: Unlocking secrets of the secreted proteins. *Proteomics* 10:799-827.
- Agrawal GK, Jwa NS, and Rakwal R (2009) Rice proteomics: end of phase I and beginning of phase II. *Proteomics* 9:935-963.
- Agrawal GK, Pedreschi R, Barkla BJ, Bindschedler LV, Cramer R, Sarkar A, Renaut J, Job D, and Rakwal R (2012a) Translational plant proteomics: A perspective. *J Proteomics* 75:4588-4601.
- Agrawal GK and Rakwal R (2006) Rice proteomics: A cornerstone for cereal food crop proteomes. *Mass Spec Rev* 25: 1-53.
- Agrawal GK and Rakwal R (2008a) *Plant Proteomics: Technologies, Strategies, and Applications*. Eds. GK Agrawal and R Rakwal, John Wiley & Sons, Inc., Hoboken, NJ, USA.
- Agrawal GK and Rakwal R (2008b) Rice proteomics at a glance. In: *Plant Proteomics: Technologies, Strategies, and Applications*. Eds. GK Agrawal and R Rakwal, John Wiley & Sons, Inc., Hoboken, NJ, USA, Chapter 11.

- Agrawal GK, Sarkar A, Agrawal R, Ndimba BK, Tanou G, Dunn MJ, Kieselbach T, Cramer R, Wienkoop S, Chen S, Rafudeen MS, Deswal R, Barkla BJ, Weckwerth W, Heazlewood JL, Renaut J, Job D, Chakraborty N, and Rakwal R (2012b) Boosting the globalization of plant proteomics through INPPO: Current developments and future prospects. *Proteomics* 12: 359-368.
- Ahsan N, Lee SH, Lee DG, Lee H, Lee SW, Bahk JD, and Lee BH (2007a) Physiological and protein profiles alternation of germinating rice seedlings exposed to acute cadmium toxicity. *C R Biol* 330: 735-746.
- Ahsan N, Lee DG, Lee SH, Kang KY, Lee JJ, Kim PJ, Yoon HS, Kim JS, and Lee BH (2007b) Excess copper induced physiological and proteomic changes in germinating rice seeds. *Chemosphere* 67: 1182-1193.
- Ahsan N, Lee DG, Alam I, Kim PJ, Lee JJ, Ahn YO, Kwak SS, Lee IJ, Bahk JD, Kang KY, Renaut J, Komatsu S, and Lee BH (2008a) Comparative proteomic study of arsenic-induced differentially expressed proteins in rice roots reveals glutathione plays a central role during As stress. *Proteomics* 8: 3561-3576.
- Ahsan N, Lee DG, Lee KW, Alam I, Lee SH, Bahk JD, and Lee BH (2008b) Glyphosate-induced oxidative stress in rice leaves revealed by proteomic approach. *Plant Physiol Biochem* 46: 1062-1070.
- Ahsan N, Lee DG, Kim KH, Alam I, Lee SH, Lee KW, Lee H, and Lee BH (2010) Analysis of arsenic stress-induced differentially expressed proteins in rice leaves by two-dimensional gel electrophoresis coupled with mass spectrometry. *Chemosphere* 78: 224-231.
- Bradshaw R and Burlingame A (2005) From proteins to proteomics. *IUBMB Life* 57: 267-272.
- Jung YH, Jeong SH, Kim SH, Singh R, Lee J, Cho YS, Agrawal GK, Rakwal R, and Jwa NS (2012) Secretome analysis of *Magnaporthe oryzae* using *in vitro* systems. *Proteomics* 12: 878-900.
- Jung YH, Lee JH, Agrawal GK, Rakwal R, Kim JA, Shim JK, Lee SK, Jeon JS, Koh HJ, Lee YH, Iwahashi H, and Jwa NS (2005) The rice (*Oryza sativa*) Blast lesion mimic mutant, blm, may confer resistance to blast pathogens by triggering multiple defense-associated signaling pathways. *Plant Physiol Biochem* 43: 397-406.
- Jung YH, Rakwal R, Agrawal GK, Shibato J, Kim JA, Lee MO, Choi PK, Jung SH, Kim SH, Koh HJ, Yonekura M, Iwahashi H, and Jwa NS (2006) Differential expression of defense/stress-related marker proteins in leaves of a unique rice blast lesion mimic mutant (blm). *J Proteome Res* 5: 2586-2598.
- Kamal AHM, Kim KH, Shin KH, Kim DE, Oh MW, Choi JS, Hirano H, Heo HY, and Woo SH (2010) Proteomics-based dissection of biotic stress responsive proteins in bread wheat (*Triticum aestivum* L.). *African J Biotechnol* 9:7239-7255.
- Kim ST, Cho KS, Jang YS, and Kang KY (2001) Two-dimensional electrophoretic analysis of rice proteins by polyethylene glycol fractionation for protein arrays. *Electrophoresis* 22: 2103-2109.
- Kim ST, Kim HS, Kim HJ, Kim SG, Kang SY, Lim DB, and Kang KY (2003a) Prefractionation of protein samples for proteome analysis by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Mol Cells* 16: 316-322.
- Kim ST, Cho KS, Yu S, Kim SG, Hong JC, Han C, Bae DW, Nam MH, and Kang KY (2003b) Proteomic analysis of differentially expressed proteins induced by rice blast fungus and elicitor in suspension-cultured rice cells. *Proteomics* 3: 2368-2378.
- Kim ST, Kim SG, Hwang DH, Kang SY, Kim HJ, Lee BH, Lee JJ, and Kang KY (2004a) Proteomic analysis of pathogen-responsive proteins from rice leaves induced by rice blast fungus, *Magnaporthe grisea*. *Proteomics* 4: 3569-3578.
- Kim ST, Yu S, Kim SG, Kim HJ, Kang SY, Hwang DH, Jang YS, and Kang KY (2004b) Proteome analysis of rice blast fungus (*Magnaporthe grisea*) proteome during appressorium formation. *Proteomics* 4: 3579-3587.
- Kim ST, Kim SG, Kang YH, Wang Y, Kim JY, Yi N, Kim JK, Rakwal R, Koh HJ, and Kang KY (2008) Proteomics analysis of rice lesion mimic mutant (*spl1*) reveals tightly localized probenazole-induced protein (PBZ1) in cells undergoing programmed cell death. *J Proteome Res* 7:1750-1760.
- Kim ST, Kang YH, Wang Y, Wu J, Park ZY, Rakwal R, Agrawal GK, Lee SY, and Kang KY (2009) Secretome analysis of differentially induced proteins in rice suspension-cultured cells triggered by rice blast fungus and elicitor. *Proteomics* 9: 1302-1313.
- Kwon YS, Ryu CM, Lee S, Park HB, Han KS, Lee JH, Lee K, Chung WS, Jeong MJ, Kim HK, and Bae DW (2010) Proteome analysis of *Arabidopsis* seedlings exposed to bacterial volatiles. *Planta* 232: 1355-1370.
- Lee BJ, Kwon SJ, Kim SK, Kim KJ, Park CJ, Kim YJ, Park OK, and Paek KH (2006) Functional study of hot pepper 26S proteasome subunit RPN7 induced by tobacco mosaic virus from nuclear proteome analysis. *Biochem Biophys Res Comm* 351: 405-411.
- Lee DG, Ahsan N, Lee SH, Kang KY, Lee JJ, and Lee BH (2007a) An approach to identify cold-induced low-abundant proteins in rice leaf. *C R Biol* 330: 215-225.
- Lee DG, Ahsan N, Lee SH, Kang KY, Bahk JD, Lee IJ, and Lee BH (2007b) A proteomic approach in analyzing heat-responsive proteins in rice leaves. *Proteomics* 7: 3369-3383.
- Lee DG, Ahsana N, Lee SH, Lee JJ, Bahk JD, Kang KY, and Lee BH (2009) Chilling stress-induced proteomic changes in rice roots. *J Plant Physiol* 166: 1-11.
- Lee K, Bae DW, Kim SH, Han HJ, Liu X, Park HC, Lim CO, Lee SY, and Chung WS (2010) Comparative proteomic analysis of the short-term responses of rice roots and leaves to cadmium. *J Plant Physiol* 167: 161-168.
- Lee Y, Bea S, Seo SG, Shim IS, Kim SH, and Kim ST (2011) Korean plant proteomics: pioneers in plant stress physiology. *J Plant Biotechnol* 38: 151-161.
- Lee YW, Shim YS, Kim YH, and Kim SM (2008) Interpretation of glufosinate reactive mechanism and proteome variation in rice (*Oryza sativa*). *Kor J Weed Sci* 28: 101-110.
- Oh IS, Park AR, Bae MS, Jae KS, Kim YS, Lee JE, Kang NY, Lee S, Cheong H, and Park OK (2005) Secretome analysis reveals an *Arabidopsis* lipase involved in defense against *Alternaria brassicicola*. *Plant Cell* 17: 2832-2847.
- Rakwal R and Agrawal GK (2003) Rice proteomics: current status and future perspectives. *Electrophoresis* 24: 3378-3389.
- Rim SO, Lee JH, Hwang SK, Suh SJ, Lee JM, Rhee IK, and Kim JG (2006) Proteome analysis of waito-c rice seedlings treated with culture fluid of gibberellin-producing fungus, *Fusarium proliferatum* KGL0401. *J Microbiol Biotechnol* 16:1990-1994.
- Ryu HS, Song MY, Kim CY, Han M, Lee SK, Ryoo N, Cho JI, Hahn TR, and Jeon JS (2009) Proteomic analysis of rice mutants susceptible to *Magnaporthe oryzae*. *Plant Biotechnol Rep* 3:167-174.

- Shin PG, Chang AC, Hong SC, Lee KS, Lee KH, and Lee YB (2008) Changes of rice storage proteins affected by dry and storage temperature. *Korean J Environ Agric* 27: 456-459.
- Swaminathan MS (2000) An evergreen revolution. *Biologist* 47: 85-89.
- Thiellement H, Zivy M, Damerval C, and Mechin V (2007) *Plant Proteomics: Methods and Protocols*. Eds. H. Thiellement, M. Zivy, C. Damerval and V. Mechin, Humana Press.
- Wang Y, Wu J, Park ZY, Kim SG, Rakwal R, Agrawal GK, Kim ST, and Kang KY (2011) Comparative Secretome Investigation of *Magnaporthe oryzae* Proteins Responsive to Nitrogen Starvation. *J Proteome Res* 10:3636-3148.
- Wehmeyer N, Hernandez LD, Finkelstein RR, and Vierling E (1996) Synthesis of small heat-shock proteins is part of the developmental program of late seed maturation. *Plant Physiol* 112: 747-757.